Hormone Foundation





CLINICAL APPLICATIONS

- Supports Favorable Estrogen Metabolism
- Supports Normal Hormone Balance
- Maintains Normal Detoxification Pathways
- Supports Gut Health for Optimal Estrogen Metabolism and Elimination
- Maintains Normal Inflammatory Balance

HORMONE HEALTH

Hormone Foundation is a comprehensive formula designed to support the body's natural hormone metabolism and detoxification. It promotes balanced estrogen pathways, maintains normal inflammatory responses, and contributes to overall wellbeing. At the core of Hormone Foundation is a carefully selected blend of ingredients that work synergistically to promote efficient hormone processing and cellular resilience. By supporting all phases of estrogen metabolism, optimizing methylation and antioxidant protection, and reinforcing gut-liver axis function, Hormone Foundation delivers targeted, multi-system support to help maintain hormonal equilibrium.

Overview

Hormone balance plays a central role in overall health and vitality for both men and women, influencing everything from energy and mood to metabolism, cognition, and long-term wellness. While hormonal fluctuations are a normal part of life, most individuals will experience periods of hormonal irregularities—whether related to life stage, stress, environment, or underlying health concerns. In women, these hormonal fluctuations can contribute to irregular menstrual cycles, fertility challenges, weight gain, fatigue, or mood swings. In men, shifts in testosterone-estrogen ratios—often due to aging, medications, body habitus, or genetic predisposition—can affect energy, mental clarity, libido, and overall vitality.

Hormone Foundation was developed to target the foundational systems that influence hormone balance, including estrogen metabolism, liver detoxification, gut health, and oxidative stress. The interconnected functions of the liver, gut, and cellular antioxidant defenses play a vital role in how hormones

are processed, transformed, and eliminated. Addressing these pathways is essential not only for managing hormonal shifts, but also for supporting the body's resilience and long-term hormonal equilibrium.

This comprehensive, systems-based approach is what makes Hormone Foundation unique: It supports the body's natural ability to process hormones efficiently, neutralize reactive metabolites, and maintain healthy estrogen balance through targeted detoxification and defense mechanisms—without overwhelming or overstimulating a single pathway. It does this by utilizing key bioactives including the Complete Turmeric Matrix, hops strobile extract, black kale extract, Quatrefolic® folate, epigallocatechin gallate (EGCg), and tributyrin (CoreBiome®).

Complete Turmeric Matrix (CTM)†

Turmeric has long been valued for its antioxidant and healthsupporting properties, with growing recognition of its role in hormone balance and the reproductive system.^{1,2} The CTM provides a full spectrum of curcuminoids and essential oils that support the body's natural detoxification processes, including the normal metabolism of estrogens through the activity of Phase I and Phase II enzymes. Research highlights turmeric's ability to maintain a balanced inflammatory response, microbiome composition, and healthy cellular function.³⁻⁶ In the context of hormone health, studies suggest that turmeric supports normal cellular processes in hormonesensitive tissues and promotes balanced estrogen metabolism through its influence on detoxification pathways.7 Its ability to enhance antioxidant protection and modulate key antioxidant regulatory proteins, such as Nrf2, further reinforces its role in overall health and wellbeing.8



Hops Strobile Extract[†]

Hops strobile extract supports the body's natural estrogen metabolism pathways by supporting Phase I detoxification and helping maintain aromatase (CYP19A1) activity within a normal range. 9,10 It aids in the natural conversion of estrogen into beneficial metabolites, supporting overall hormonal balance. Hops contains bioactive compounds, including xanthohumol and 8-prenylnaringenin (8-PN), which help maintain antioxidant activity and a balanced inflammatory response, in addition to promoting cellular health. 11,12 Research suggests that the phytoestrogenic properties of hops contribute to its ability to support menopausal comfort, as clinical trials have observed improvements in hot flashes when compared to a placebo. 13,14

Additionally, hops constituents have been studied for their role in estrogen metabolism. Xanthohumol has been shown to influence detoxification enzymes, promote oxidative balance through Nrf2 activation, and support normal inflammatory pathways. ¹² Other compounds, such as 6-prenylnaringenin (6-PN), have been researched for their ability to upregulate specific estrogen detoxification enzymes. ¹⁵ Prenylated phenols in hops also contribute to antioxidant support by promoting glutathione synthesis and other antioxidative enzymes. ^{12,16}

Black Kale Extract (18% Glucoraphanin)†

Black kale extract is a rich source of glucoraphanin, a naturally occurring, sulfur-containing compound that supports the body's normal detoxification pathways. Glucoraphanin is a precursor to sulforaphane, which has been shown to promote Phase II detoxification enzyme activity, assisting in the efficient processing and elimination of endogenous and environmental compounds. By supporting sulfation pathways, black kale extract helps maintain balanced hormone metabolism and overall cellular resilience. Research indicates that glucoraphanin contributes to normal antioxidant defenses, further promoting cellular health and physiological equilibrium. B. 19

Folate (as Quatrefolic® 5-MTHF)†

Folate (as Quatrefolic®) is essential for methylation, a key biochemical process involved in estrogen metabolism. It facilitates the conversion of catechol estrogens, such as 2-hydroxyestrone and 4-hydroxyestrone, into methylated derivatives for safer elimination, thereby promoting hormonal balance and reducing oxidative stress. Quatrefolic®, a bioavailable form of 5-MTHF, ensures adequate methylation capacity to support estrogen detoxification. Additionally, folate plays a critical role in DNA synthesis and repair, helping to protect cells from oxidative stress associated with estrogen metabolism. ^{20,21}

Epigallocatechin Gallate (EGCg)[†]

EGCg, a green tea polyphenol, enhances antioxidant activity, which helps maintain cellular integrity during the natural processing of estrogen. EGCg can also influence estrogen receptor (ER)-dependent expression of estrogen target genes and ER-independent signaling pathways.²² This is demonstrated in a study of urinary estrogen profiles in green tea drinkers that showed a favorable modification of estrogen metabolism with a reduction of 16-OH estrogens as well as increased ratio of methylated 2-OH to 4-OH catechol estrogens, a process associated with maintaining hormonal equilibrium.²³

Tributyrin (CoreBiome®)†

Tributyrin is a precursor to butyrate, a short-chain fatty acid that supports a healthy digestive environment and plays a key role in estrogen metabolism. As a bioavailable form of butyrate, tributyrin promotes normal gut barrier function, helping to maintain balanced microbial beta-glucuronidase activity and supporting the gut-liver axis in hormone detoxification. ^{24,25,26} By strengthening the gut barrier, butyrate reduces the reabsorption of estrogen metabolites that have already been processed by the liver. ²⁷ Through these actions, tributyrin supports efficient estrogen elimination and helps maintain normal hormonal balance.



Directions

1 capsule per day or as recommended by your healthcare professional.

Does Not Contain

Gluten, yeast, artificial colors or flavors.

Caution

If you are pregnant or nursing, consult your physician before taking this product.

Supplement Serving Size 1 Capsule	nt Fac	cts
Servings Per Container 30 & 60		
	Amount Per Serving	% Daily Value
Calories	5	
Total Fat	0.5 g	<1%*
Saturated Fat	0.5 g	3%*
Folate (from 235 mcg as Quatrefolic® (6S)-5-Methyltetrahydrofolic acid glucos	400 mcg DFE samine salt)	100%
Tributyrin (as CoreBiome®)	300 mg	*
Black Kale Seed Extract (Standardized to contain 18 mg Glucora	100 mg aphanin)	*
Turmeric Root Extract (Complete Turmeric Matrix) (Standardiz 45-55% Curcuminoids, 3-8% Volatile O		*
Green Tea Leaf Extract (Standardized to contain 45% EGCg (E	50 mg pigallocatechin gallat	* te))
Hops Extract	50 mg	*
* Daily Value not established.		

Other Ingredients: Hypromellose (Natural Vegetable Capsule), Medium Chain Triglycerides, Candelilla Wax and Sunflower Lecithin.

ID# 166030 30 Capsules ID# 166060 60 Capsules

References

- Kamal DAM, Salamt N, Yusuf ANM, Kashim MIAM, Mokhtar MH. Potential health benefits of curcumin on female reproductive disorders: A review. *Nutrients* 2021;13(9):3126. doi:10.3390/nu13093126.
- 2. Sirotkin AV. The influence of turmeric and curcumin on female reproductive processes. *Planta Med* 2022;88(12):1020-1025. doi:10.1055/a-1542-8992.
- 3. Jiang TA. Health benefits of culinary herbs and spices. *JAOAC Int* 2019;102(2):395-411. doi:10.5740/jaoacint.18-0418.
- 4. Hewlings SJ, Kalman DS. Curcumin: A review of its effects on human health. *Foods* 2017;6(10):92. doi:10.3390/foods6100092.

- 5. Shah M, Murad W, Mubin S, Ullah O, Rehman NU, Rahman MH. Multiple health benefits of curcumin and its therapeutic potential. *Environ Sci Pollut Res Int* 2022;29(29):43732-43744. doi:10.1007/s11356-022-20137-w.
- Liu S, Liu J, He L, et al. A comprehensive review on the benefits and problems of curcumin with respect to human health. *Molecules* 2022;27(14):4400. doi:10.3390/ molecules27144400.
- 7. Hodges RE, Minich DM. Modulation of metabolic detoxification pathways using foods and food-derived components: A scientific review with clinical application. *J Nutr Metab* 2015;2015:760689. doi:10.1155/2015/760689. PMID: 26167297; PMCID: PMC4488002.
- 8. Bak MJ, Das Gupta S, Wahler J, Suh N. Role of dietary bioactive natural products in estrogen receptor-positive breast cancer. *Semin Cancer Biol* 2016;40-41:170-191. doi:10.1016/j.semcancer.2016.03.001.
- 9. Monteiro R, Becker H, Azevedo I, Calhau C. Effect of hop (Humulus lupulus L.) flavonoids on aromatase (estrogen synthase) activity. *J Agric Food Chem* 2006;54(8):2938-2943. doi:10.1021/jf053162t.
- 10. Monteiro R, Faria A, Azevedo I, Calhau C. Modulation of breast cancer cell survival by aromatase-inhibiting hop (Humulus lupulus L.) flavonoids. *J Steroid Biochem Mol Biol* 2007;105(1-5):124-130. doi:10.1016/j.jsbmb.2006.11.026.
- 11. Dostálek P, Karabín M, Jelínek L. Hop phytochemicals and their potential role in metabolic syndrome prevention and therapy. *Molecules* 2017;22(10):1761. doi:10.3390/molecules22101761.
- 12. Bolton JL, Dunlap TL, Hajirahimkhan A, et al. The multiple biological targets of hops and bioactive compounds. *Chem Res Toxicol* 2019;32(2):222-233. doi:10.1021/acs. chemrestox.8b00345. Epub 2019 Jan 22. Erratum in: *Chem Res Toxicol* 2019;32(8):1732. PMID: 30608650; PMCID: PMC6643004.
- 13. Aghamiri V, Mirghafourvand M, Mohammad-Alizadeh-Charandabi S, Nazemiyeh H. The effect of hop (Humulus lupulus L.) on early menopausal symptoms and hot flashes: A randomized placebo-controlled trial. *Complement Ther Clin Pract* 2016;23:130-135. doi:10.1016/j.ctcp.2015.05.001.
- 14. Keiler AM, Zierau O, Kretzschmar G. Hop extracts and hop substances in the treatment of menopausal complaints. *PlantaMed* 2013;79(7):576-579. doi:10.1055/s-0032-1328330.



- 15. Wang S, Dunlap TL, Howell CE, et al. Hop (Humulus lupulus L.) extract and 6-prenylnaringenin induce P450 1A1 catalyzed estrogen 2-hydroxylation. *Chem Res Toxicol* 2016;29(7):1142-1150. doi:10.1021/acs. chemrestox.6b00112.
- 16. Hitzman RT, Dunlap TL, Howell CE, et al. 6-Prenylnaringenin from hops disrupts ERα-mediated downregulation of CYP1A1 to facilitate estrogen detoxification. *Chem Res Toxicol* 2020;33(11):2793-2803. doi:10.1021/acs. chemrestox.0c00194.
- 17. Fahey JW, Talalay P. Antioxidant functions of sulforaphane: A potent inducer of Phase II detoxification enzymes. *Food Chem Toxicol* 1999;37(9-10):973-979. doi:10.1016/s0278-6915(99)00082-4.
- 18. Dinkova-Kostova AT, Kostov RV. Glucosinolates and isothiocyanates in health and disease. *Trends Mol Med* 2012;18(6):337-347. doi:10.1016/j.molmed.2012.04.003.
- 19. Kaufman-Szymczyk A, Majewski G, Lubecka-Pietruszewska K, Fabianowska-Majewska K. The role of sulforaphane in epigenetic mechanisms, including interdependence between histone modification and DNA methylation. *Int J Mol Sci* 2015;16(12):29732-29743. doi:10.3390/ijms161226194.
- 20. Goodman J, Lavigne JA, Wu K, et al. COMT genotype, micronutrients in the folate metabolic pathway, and breast cancer risk. *Carcinogenesis* 2001;22(10):1661-1665. doi:10.1093/carcin/22.10.1661.
- 21. National Institutes of Health. Office of Dietary Supplements Folate. NIH.gov. Published 2017. Available from: https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/.
- 22. Bak MJ, Das Gupta S, Wahler J, Suh N. Role of dietary bioactive natural products in estrogen receptor-positive breast cancer. *Semin Cancer Biol* 2016;40-41:170-191. doi:10.1016/j. semcancer.2016.03.001.
- 23. Fuhrman BJ, Pfeiffer RM, Wu AH, et al. Green tea intake is associated with urinary estrogen profiles in Japanese-American women. *Nutr J* 2013;12:25. doi:10.1186/1475-2891-12-25.
- 24. Martin-Gallausiaux C, Marinelli L, Blottière HM, Larraufie P, Lapaque N. SCFA: Mechanisms and functional importance in the gut. *Proc Nutr Soc* 2021;80:37-49. doi:10.1017/S0029665120006916.

- 25. Yang N, Lan T, Han Y, et al. Tributyrin alleviates gut microbiota dysbiosis to repair intestinal damage in antibiotic-treated mice. *PLoS One* 2023;18(7):e0289364. doi:10.1371/journal. pone.0289364.
- 26. Ervin SM, Li H, Lim L, et al. Gut microbial β -glucuronidases reactivate estrogens as components of the estrobolome. *J Biol Chem* 2019;294(47):18586-18599. doi:10.1074/jbc. RA119.010950.
- 27. Flores R, Shi J, Fuhrman B, et al. Fecal microbial determinants of fecal and systemic estrogens and estrogen metabolites: A cross-sectional study. *J Transl Med* 2012;10:253. doi:10.1186/1479-5876-10-253.

